

Spinning strings, cosmic dislocations, and chronology protection

V. A. De Lorenci* and E. S. Moreira, Jr.†

*Instituto de Ciências Exatas, Universidade Federal de Itajubá,
Av. BPS 1303 Pinheirinho, 37500-903 Itajubá, Minas Gerais, Brazil*

(Dated: September, 2003)

A massless scalar field is quantized in the background of a spinning string with cosmic dislocation. By increasing the spin density toward the dislocation parameter, a region containing closed timelike curves (CTCs) eventually forms around the defect. Correspondingly, the propagator tends to the ordinary cosmic string propagator, leading therefore to a mean-square field fluctuation, which remains well behaved throughout the process, unlike the vacuum expectation value of the energy-momentum tensor, which diverges due to a subtle mechanism. These results suggest that back reaction leads to the formation of a “horizon” that protects from the appearance of CTCs.

PACS numbers: 04.62.+v, 04.20.Gz, 11.27.+d

Investigations on quantum theory around spinning defects go back to the late 1980s with the study of quantum mechanics of relativistic particles on the spinning cone [1]. Such a background is the Kerr-like solution of the Einstein equations in three dimensions, whose line element is given by (throughout the text $c = \hbar = 1$, and the metric parameters are nonnegative)

$$ds^2 = (d\tau + Sd\theta)^2 - dr^2 - \alpha^2 r^2 d\theta^2, \quad (1)$$

where S and α are the spin and the disclination parameter, respectively [2]. Clearly Minkowski spacetime corresponds to $S = 0$ and $\alpha = 1$. Lifting the geometry in Eq. (1) to four dimensions, one obtains the gravitational background around a spinning cosmic string [3], for which

$$ds^2 = (d\tau + Sd\theta)^2 - dr^2 - \alpha^2 r^2 d\theta^2 - d\xi^2. \quad (2)$$

An inspection of Eqs. (1) and (2) shows that the region for which $r < S/\alpha$ contains CTCs, resulting that when $S \neq 0$ the corresponding spacetimes are not globally hyperbolic. It is not clear if quantum theory makes sense in nonglobally hyperbolic spacetimes [4]. In fact, quantum mechanics on the spinning cone has shown that $S \neq 0$ spoils unitarity [1]. In the context of the second quantization around spinning cosmic strings [5], a recent analysis has revealed that a nonvanishing spin density S leads to divergent vacuum fluctuations [6].

In order to recover boost invariance along the symmetry axis, the authors in Ref. [7] have “amended” the geometry in Eq. (2) by postulating a cosmic dislocation, such that

$$ds^2 = (d\tau + Sd\theta)^2 - dr^2 - \alpha^2 r^2 d\theta^2 - (d\xi + \kappa d\theta)^2, \quad (3)$$

whose metric tensor fits as solution of the Einstein equations, as well as solution of the Einstein-Cartan equations [8, 9]. When $S > \kappa$, the region for which $r < \sqrt{S^2 - \kappa^2}/\alpha$ contains CTCs. When $S < \kappa$ though, the spacetime is globally hyperbolic.

Vacuum fluctuations typically diverge on the Cauchy surface (chronology horizon), which separates a region with CTCs from another without CTCs (for a review see Ref. [10]). This fact has led to the chronology protection conjecture, according to which, physical laws do not allow the appearance of CTCs (“time machines”) [11]. Although the geometry in Eq. (3) does not contain any Cauchy horizon [for $S > \kappa$, Eq. (3) describes an “eternal time machine”], it might be clarifying to study quantum effects in the corresponding spacetime as the metric parameters are adjusted such that CTCs are about to form. Using a massless scalar field as a probe, this work implements such an investigation by considering $S < \kappa$ and by taking $S \rightarrow \kappa$, i.e., arbitrarily close to the point when the spacetime is about to become nonglobally hyperbolic.

According to Ref. [7], when the metric parameters in Eq. (3) satisfy $S < \kappa$, a Lorentz frame exists with respect to which the spin density vanishes (and one might say, in this case, that $S \neq 0$ is a kinematic effect). Indeed, by performing the following Lorentz transformation in the $\tau - \xi$ plane,

$$t = \frac{\tau - v\xi}{\sqrt{1 - v^2}} \quad z = \frac{\xi - v\tau}{\sqrt{1 - v^2}} \quad v := S/\kappa, \quad (4)$$

Eq. (3) can be recast as

$$ds^2 = dt^2 - dr^2 - \alpha^2 r^2 d\theta^2 - (dz + \kappa' d\theta)^2, \quad (5)$$

describing the gravitational background of a cosmic dislocation with dislocation parameter

$$\kappa' := \sqrt{\kappa^2 - S^2}, \quad (6)$$

and for which the usual identification

$$(t, r, \theta, z) \sim (t, r, \theta + 2\pi, z) \quad (7)$$

is observed. It should be noted that the background of a spinning string can only be seen as that of a cosmic dislocation when $S < \kappa$, since when $S \geq \kappa$ the boost in Eq. (4) becomes singular.

Vacuum fluctuations of a massless scalar field ϕ around a cosmic dislocation have recently been reported in the literature [12]. Equations (5) and (7) show that when $\kappa' \rightarrow 0$ the corresponding vacuum fluctuations become those associated with an ordinary cosmic string (see, e.g., [13]). It follows that all scalar vacuum averages observed from the frame corresponding to Eq. (3) are meant to remain finite as S is taken arbitrarily close to the critical value κ . For example, as $S \rightarrow \kappa$ the mean-square field fluctuation approaches

$$\langle \phi^2(r) \rangle = \frac{1}{48\pi^2 r^2} (\alpha^{-2} - 1), \quad (8)$$

which is finite (away from the defect).

Turning to the vacuum expectation value of the energy-momentum tensor $\langle T^\mu_\nu \rangle$, it is more convenient to use local inertial coordinates (T, r, φ, Ξ) and (t, r, φ, Z) associated with Eqs. (3) and (5), respectively, which are defined as $T := \tau + S\theta$, $\varphi := \alpha\theta$, $\Xi := \xi + \kappa\theta$ and $Z := z + \kappa'\theta$. In terms of these coordinates, both Eqs. (3) and (5) become the Minkowski line element written in cylindrical coordinates, and Eq. (7) leads to $(t, r, \varphi, Z) \sim (t, r, \varphi + 2\pi\alpha, Z + 2\pi\kappa')$, revealing a “space-helical” structure. [Representing the spinning string by a rotating helix, Eq. (4) leads to the frame travelling through the symmetry axis, and for which the helix does not rotate]. It should be mentioned that when the coordinates (T, r, φ, Ξ) are used, the background appears to have also a “time-helical” structure [2].

Using Eq. (4), one finds out that the energy density $\langle T^T_T \rangle$ in the spinning string inertial frame (T, r, φ, Ξ) is related with $\langle T^\mu_\nu \rangle$ in the cosmic dislocation inertial frame (t, r, φ, Z) by

$$\langle T^T_T \rangle = \frac{\langle T^t_t \rangle - v^2 \langle T^Z_Z \rangle}{1 - v^2}. \quad (9)$$

A superficial investigation may suggest that the relativistic factor in Eq. (9) will make $\langle T^T_T \rangle$ to diverge as $S \rightarrow \kappa$ [$v \rightarrow 1$, cf. Eq. (4)]. However, as will be seen shortly, this is incorrect — $\langle T^T_T \rangle$ indeed diverges as $S \rightarrow \kappa$; but the mechanism through which that operates is rather subtle and does not involve any relativistic factor. Another pitfall consists of carrying over Eq. (9) the fact that $S \rightarrow \kappa$ [$\kappa' \rightarrow 0$, cf. Eq. (6)] leads to $\langle T^Z_Z \rangle \rightarrow \langle T^t_t \rangle$ [12], and then to conclude (incorrectly) that $\langle T^T_T \rangle \rightarrow \langle T^t_t \rangle$. The flaw in this argument will be clear in the following.

At this point one recalls that $\langle T^t_t \rangle$ and $\langle T^Z_Z \rangle$ in Eq. (9) are vacuum fluctuations in the background of a cosmic dislocation with dislocation parameter κ' . As is explained in Ref. [12], $\langle T^t_t \rangle$ and $\langle T^Z_Z \rangle$ can be obtained by letting a certain differential operator to act on the corresponding renormalized propagator $D^{(\alpha, \kappa')}(x, \bar{x})$, according to the prescription in Eq. (17) of Ref. [12]. Proceeding along these lines, it follows that

$$\langle T^Z_Z \rangle - \langle T^t_t \rangle = -i \lim_{\bar{x} \rightarrow x} (\partial_t \partial_{\bar{t}} + \partial_Z \partial_{\bar{Z}}) D^{(\alpha, \kappa')}(x, \bar{x}), \quad (10)$$

and by letting the derivatives to act on the expression of $D^{(\alpha, \kappa')}(x, \bar{x})$ in Eq. (12) of Ref. [12], Eq. (10) yields

$$\langle T^Z_Z \rangle = \langle T^t_t \rangle + \frac{\kappa'^2}{r^6} f_\alpha(\kappa'^2/r^2), \quad (11)$$

where

$$f_\alpha(x) := -\frac{1}{2} \int_0^\infty d\lambda \sum_{n=1}^\infty \frac{n^2 [\lambda^2 - \pi^2(4\alpha^2 n^2 - 1)]}{[\pi^2(2\alpha n + 1)^2 + \lambda^2] [\pi^2(2\alpha n - 1)^2 + \lambda^2] [\cosh^2(\lambda/2) + n^2 \pi^2 x]^3}. \quad (12)$$

A quick power counting in Eq. (12) gives that $f_\alpha(x)$ diverges at $x = 0$ (when α is finite). A more careful analysis shows that $f_\alpha(x)$ diverges as $x \rightarrow 0$; but it does so mildly since $x f_\alpha(x) \rightarrow 0$.

By inserting Eq. (11) in Eq. (9) and recalling that $\kappa'^2 = \kappa^2(1 - v^2)$, one ends up with

$$\langle \mathcal{T}^T_T \rangle = \langle T^t_t \rangle - \frac{S^2}{r^6} f_\alpha (\kappa'^2/r^2). \quad (13)$$

Considering $S \rightarrow \kappa$ in Eq. (13), $\langle T^t_t \rangle$ approaches the ordinary cosmic string expression (and therefore remains finite), whereas the term carrying f_α diverges [one sees that the flaw mentioned above consists in manipulating improperly the numerator in Eq. (9) when κ' is very small]. One might say that $S = \kappa$ plays the role of a chronology horizon where mechanisms of chronology protection are expected to take place. The divergence in $\langle \mathcal{T}^T_T \rangle$ confirms this expectation. For completeness, the other components of $\langle \mathcal{T}^\mu_\nu \rangle$ are displayed below

$$\langle \mathcal{T}^\mu_\nu \rangle = \begin{pmatrix} \langle T^t_t \rangle - (S^2/r^6)f_\alpha & 0 & (S/\kappa') \langle T^Z_\varphi \rangle & (\kappa S/r^6)f_\alpha \\ 0 & \langle T^r_r \rangle & 0 & 0 \\ -(S/\kappa') \langle T^\varphi_Z \rangle & 0 & \langle T^\varphi_\varphi \rangle & (\kappa/\kappa') \langle T^\varphi_Z \rangle \\ -(\kappa S/r^6)f_\alpha & 0 & (\kappa/\kappa') \langle T^Z_\varphi \rangle & \langle T^Z_Z \rangle + (S^2/r^6)f_\alpha \end{pmatrix}, \quad (14)$$

with f_α evaluated at κ'^2/r^2 . The approximate behavior of $\langle \mathcal{T}^\mu_\nu \rangle$ as $S \rightarrow \kappa$, at a given distance r from the defect, can be obtained from Eq. (14) by considering the expressions for $\langle T^\mu_\nu \rangle$ in Ref. [12]. For example, if ϕ is conformally coupled, it follows that

$$\langle \mathcal{T}^\mu_\nu \rangle = \frac{1}{r^4} \begin{pmatrix} -(S^2/r^2)f_\alpha - A & 0 & SB & (\kappa S/r^2)f_\alpha \\ 0 & -A & 0 & 0 \\ -SB/r^2 & 0 & 3A & \kappa B/r^2 \\ -(\kappa S/r^2)f_\alpha & 0 & \kappa B & (S^2/r^2)f_\alpha - A \end{pmatrix}, \quad (15)$$

where $A(\alpha) := (\alpha^{-4} - 1)/1440\pi^2$ and $B(\alpha)$ is defined as in Eq. (20) of Ref. [12] [$B(\alpha = 1) = 1/60\pi^2$]. Clearly the components that diverge as $S \rightarrow \kappa$ are those containing f_α . At this point it is pertinent to note that, as mentioned previously, $\langle \phi^2 \rangle$ remains well behaved as S approaches the critical value κ [see Eq. (8)]. If $f_\alpha(x)$ were not divergent at $x = 0$, $\langle \mathcal{T}^\mu_\nu \rangle$ would also remain finite as $S \rightarrow \kappa$ (and that would suggest violation of chronology protection).

The following are simple facts that help to figure out the physical implication of the results reported above. One begins by taking S infinitesimally smaller than κ , say $S = \kappa - \delta$. Then, a finite interval of proper time $\Delta\tau$ as measured in the spinning string frame [cf., Eq. (3)] would appear an arbitrarily large interval of time measured in the corresponding cosmic dislocation frame [cf., Eq. (5)], since the latter would be traveling nearly at the speed of light (recall that $v = S/\kappa$). If $\Delta\tau$ is the interval of proper time immediately before the emergence of a region containing CTCs (which corresponds to $\delta = 0$), it follows that such an event never would be detected in the cosmic dislocation frame. Moreover, as $\delta \rightarrow 0$, the semiclassical metric tensor in the cosmic dislocation frame approaches that found in Ref. [14], which presents mild contributions from back reaction, as is typically the case. Considering the divergences in the vacuum expectation value of the energy-momentum tensor, such mild contributions (and that is a crucial point) would be hugely amplified in the spinning string frame, suggesting the formation of a “horizon” when $\delta = 0$. Such a “horizon” would eventually prevent the appearance of CTCs around the spinning string.

The authors are indebted to Renato Klippert for useful discussions. This work was supported by the Brazilian research agencies CNPq and FAPEMIG.

* Electronic address: delorenci@unifei.edu.br

† Electronic address: moreira@unifei.edu.br

- [1] P. S. Gerbert and R. Jackiw, Commun. Math. Phys. **124**, 229 (1989).
- [2] S. Deser, R. Jackiw, and G. 't Hooft, Ann. Phys. **152**, 220 (1984).
- [3] P. O. Mazur, Phys. Rev. Lett. **57**, 929 (1986).
- [4] S. A. Fulling, *Aspects of Quantum Field Theory in Curved Space-Time* (Cambridge University Press, Cambridge, England, 1989).
- [5] G. E. A. Matsas, Phys. Rev. D **42**, 2927 (1990).
- [6] V. A. De Lorenci and E. S. Moreira, Jr., Phys. Rev. D **63**, 027501 (2001).
- [7] D. V. Gal'tsov and P. S. Letelier, Phys. Rev. D **47**, 4273 (1993).
- [8] K. P. Tod, Class. Quantum Grav. **11**, 1331 (1994).
- [9] P. S. Letelier, Class. Quantum Grav. **12**, 471 (1995).
- [10] M. Visser, gr-qc/0204022 (unpublished).

- [11] S. W. Hawking, Phys. Rev. D **46**, 603 (1992).
- [12] V. A. De Lorenci and E. S. Moreira, Jr., Phys. Rev. D **67**, 124002 (2003).
- [13] A. G. Smith, in *The Formation and Evolution of Cosmic Strings*, Proceedings of the Cambridge Workshop, Cambridge, England, edited by G. W. Gibbons, S. W. Hawking, and T. Vachaspati (Cambridge University Press, Cambridge, England, 1990).
- [14] W. A. Hiscock, Phys. Lett. B **188**, 317 (1987).